

Tilburg University

Atypical visual-auditory predictive coding in Autism Spectrum Disorder

van Laarhoven, Thijs; Stekelenburg, Jeroen; Eussen, M.; Vroomen, Jean

Publication date:
2019

Document Version
Publisher's PDF, also known as Version of record

[Link to publication in Tilburg University Research Portal](#)

Citation for published version (APA):

van Laarhoven, T., Stekelenburg, J., Eussen, M., & Vroomen, J. (2019). *Atypical visual-auditory predictive coding in Autism Spectrum Disorder: Electrophysiological evidence from stimulus omissions*. Poster session presented at NVP Winter Conference, Egmond aan Zee, Netherlands.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Atypical Visual-Auditory Predictive Coding in Autism Spectrum Disorder

Electrophysiological Evidence from Stimulus Omissions



PRESENTER

Thijs van Laarhoven

t.j.t.m.vanlaarhoven@tilburguniversity.edu

SUMMARY

Autism Spectrum Disorder (ASD) is a pervasive neurodevelopmental disorder that has been linked to a range of perceptual processing alterations, including hypo- and hyperresponsiveness to sensory stimulation.

Recent evidence suggests that these symptoms might be related to a decreased ability to anticipate upcoming sensory stimulation.¹ The results of this study showed that individuals with ASD may be unable to fully anticipate the sensory consequences of their own motor actions.

This raised the question if the ability to predict the actions of other individuals is altered as well in ASD.

Here, we used a stimulus omission paradigm to examine the electrophysiological markers of prediction error² in auditory prediction by vision in individuals with ASD to assess their ability to anticipate the sensory consequences of others' actions.

Unexpected auditory omissions in a sequence of audiovisual recordings in which the visual motion reliably predicted the timing and content of the sound elicited **larger prediction errors** in our sample of individuals with ASD.

The current data suggest that individuals with ASD may have impairments in the ability to anticipate the sensory consequences of others' actions and support the notion that sensory prediction might be overly precise and inflexible in ASD.³


1. van Laarhoven, T., Stekelenburg, J. J., Eussen, M. L. J. M. J. M., & Vroomen, J. (2019). Electrophysiological alterations in motor-auditory predictive coding in autism spectrum disorder. *Autism Research*, 12(4), 589–599. <https://doi.org/10.1002/aur.2087>

2. Friston, K. (2005). A theory of cortical responses. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 360(1456), 815–836. <https://doi.org/10.1098/rstb.2005.1622>

3. van de Cruys, S., Evers, K., van der Hallen, R., van Eylen, L., Boets, B., de-Wit, L., & Wagemans, J. (2014). Precise minds in uncertain worlds: Predictive coding in autism. *Psychological Review*, 121(4), 649–675. <https://doi.org/10.1037/a0037665>

Violations of visual-to-auditory predictions induce larger prediction errors in Autism Spectrum Disorder

Sound Omission ERPs (measured in silence)




Take a picture to download the abstract (manuscript currently under review)

PARTICIPANTS

Group	Sample size	No. females	Mean age	Mean IQ
ASD	29	8	18.64	103.03
Typical Development	29	6	18.93	112.07

EXPERIMENTAL CONDITIONS



Visual-auditory (VA)

- video + sound
- 1232 trials (88%)

VA trials were interspersed with **168 unpredictable sound omission trials (12% of all 1400 trials)**


Visual (V)

- video only
- 200 trials

V trials were included to correct for visual activity in the sound omission trials of the VA condition (VA-V)

RESULTS

Scalp Potential Maps of Sound Omission Trials



Thijs van Laarhoven^a Jeroen J. Stekelenburg^a
Mart Eussen^b Jean Vroomen^a

a. Department of Cognitive Neuropsychology, Tilburg University, Tilburg, The Netherlands
b. Department of Autism, Yulius Mental Health Organization, Dordrecht, The Netherlands

